

TriHealth Digestive Institute

MoviPrep Colonoscopy with Fecal Transplant Prep Split Dosing with Local IV or MAC Sedation

Location of Procedure: {endo locations:17161}	
DATE: ***	Arrival Time: ***
PHYSICIAN: {tdi providers:16561}	Procedure Time: ***

Please read these entire instructions several days before your actual procedure appointment.

PRIOR TO PROCEDURE:

YOU WILL NEED TO PURCHASE:

1 MoviPrep Bowel Kit (Prescription Required. Sent to your Pharmacy @PHARM@)

Diaper Rash Cream- optional, over the counter (No prescription required)

Imodium (Over the Counter)

If you have any question, please call our office during regular business at 513-794-5600 for Ohio and 812-537-5558 for Indiana. You will be at the facility approximately 2-4 hours. Please make sure you arrange for someone to bring you to the procedure and most importantly, drive you home from your procedure. If you don't have a driver your procedure will be cancelled. There must be someone responsible for your welfare such as a friend or family member. You will not be allowed to leave the facility ALONE in a cab.

All current medications that you are taking should be continued in routine dosage unless you are taking prescribed blood thinners, iron, diabetic medications, Vitamin E or Fish Oil. If applicable, your specific instructions should be indicated below if you are on any of these types of medications. Please note that you should not stop taking your blood thinners until you get the OK with instructions from your doctor.

In addition, if you are taking Aspirin prescribed by your doctor then you should continue to take it. If you are taking Fish Oil or Vitamin E please stop taking 7 days prior to your procedure. If you take a diuretic/waterpill, don't take it the morning of your procedure.

Your Specific Medication Instructions:

You are on anticoagulation medication: {prep coag:18837}
You are on diabetic medication: {diabetic:18797}
You are on Iron Medication: {iron for preps:18794}

PLEASE DRINK PLENTY OF LIQUIDS.

2 DAYS PRIOR TO PROCEDURE

NO FRESH FRUITS, RAW VEGETABLES, CORN, SEEDS OR NUTS.

Stop all antibiotics and probiotics being used to treat C. Diff 48 hours prior to procedure.

1 DAY BEFORE PROCEDURE

Begin a **clear, liquid diet** in the morning. Drink at least 1/2 gallon of fluids throughout the day and evening **IN ADDITION** to the MoviPrep solution.

A clear, liquid diet consists of any clear fruit juice (without pulp), soft drinks, clear broth (beef or chicken bouillon is fine), coffee, tea, Kool-Aid, Gatorade, Propel water, Jello, Gelatin, regular popsicles, any type of hard candy (but no soft center or chocolate). **Do not consume dairy products or liquids with RED, BLUE, OR PURPLE COLORING ADDED.** It is recommended that you drink 32 oz. Of Gatorade.

At 6 p.m., empty 1 Pouch A and 1 Pouch B into the disposable container. Add lukewarm drinking water to the top line of the container and mix to dissolve. If preferred, mix solution ahead of time and refrigerate prior to drinking. The solution should be used within 24 hours. It might help to use a straw.

The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz.) until the container is empty. Over the next hour drink an additional 16 oz of clear liquid of your choice.

DAY OF PROCEDURE

4 hours prior to your arrival time, empty 1 Pouch A and 1 Pouch B into the disposable container. Add lukewarm drinking water to the top line of the container and mix to dissolve. If preferred, mix solution ahead of time and refrigerate prior to drinking. The solution should be used within 24 hours. It might help to use a straw.

The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz.) until the container is empty. Over the next hour drink an additional 16 oz of clear liquid of your choice.

After you finish the last 16 oz. Container of clear liquid, you are to have nothing to eat or drink until after the colonoscopy. Although this time period may be during the middle of the night, studies have shown this to be the best method to ensure a clean prep.

2 hours prior to the procedure take 4 mg Imodium with a small sip of water.

FINAL INSTRUCTIONS

More detailed instructions can be found by visiting our website: <http://www.trihealth.com/institutes-and-services/trihealth-digestive-institute/patient-information/>

- This exam is most successful if these instructions are followed exactly as stated above.
- If you use a CPAP machine at night for sleep apnea, please bring this with you.
- Bring a list of your current medications (including dosages) with you.
- Verify any questions about co-pays with your insurance company.

**If you are unable to keep this appointment you must give 72 hours (3 business days) notice to allow us to fill this appointment time with another patient or you may be subject to a \$75.00 no show charge which is not covered by your insurance.

There may be times when we leave Personal Health Information on your answering machine. If you do not wish for us to do this, please call the office at 513-794-5600 (Ohio) or 812-537-5558 (Indiana).

As is the case with any recommendations from your physician, please follow the exact instructions in taking this prep for your procedure. Doing so will help ensure your colon will be clear of stool. If the colon prep isn't finished or taken per the instructions it could lead to ineffective fecal transplant and the procedure may need to be scheduled again and the bowel prep repeated.

Please contact your insurance company for specific information around the processing of your upcoming claim. There are multiple guidelines and interpretations from each insurance company that could make your claim be applied to your deductible especially around colonoscopies. When calling your insurance company, please be clear to indicate to them you are having a colonoscopy with fecal transplant. Based on the information you have provided to us while scheduling your procedure, we will most likely use the procedure code and diagnosis listed below. Please note that these codes could potentially change based on what the physician finds during the procedure or if you indicate something different at the time of your procedure to the nurses or the physician. Once your claim has been billed we aren't able to go back and change diagnoses.

Procedure code: Colonoscopy CPT CODE: 45378 **Diagnosis code:** C. Diff ICD10 A04.7 (ICD9 008.45)

CLEAR LIQUID DIET

A liquid is "clear" if you can read something through it.

Example: When placing a glass with Apple Juice over a newspaper you are able to see the words, this is a clear liquid.

Drink plenty of fluids all day. Water can get boring, so keep a variety of clear liquids on hand.

Clear liquids include any of the following:

- Water
- White Cranberry, White Grape, Apple Juices (without any pulp)
- Honey
- Any powdered drink mix with water (Crystal Light, Kool-Aid, Etc. Except for No **RED, BLUE, OR PURPLE**)
- Any Soda (Except No **RED, BLUE, OR PURPLE**)
- Lemonade
- Sports Drinks (Except No **RED, BLUE, OR PURPLE**)
- Flavored gelatin/popsicles (No **RED, BLUE, OR PURPLE**)
- Broth/bouillon
- Coffee/tea with sugar (No Milk or Creamer)
- Gum
- Hard Candies (Do not chew)
- Special K Protein Water
- Although alcohol is a clear liquid, it can make you dehydrated. You should **NOT** drink alcohol during the preparation for your test.

Sample Menus of Clear Liquid Diets (These are only samples you do not have to use these)

Menu #1

Breakfast: Hot tea with lemon juice and 1 tsp sugar (no pulp); Apple juice (8 oz); Gelatin (1 cup)

Lunch: Hot tea with lemon (no pulp) and sugar; White Grape Juice (8 oz); Fruit Ice/popsicles (1 cup); Broth (8 oz)

Snack: Fruit juice (apple, white cranberry or white grape, 8 oz.); Gelatin (1 cup)

Dinner: Hot tea with lemon (no pulp) and sugar; Apple juice (8 oz); Broth (8 oz); Fruit ice/popsicles (1 cup)

Menu #2

Breakfast: 1 glass fruit juice; 1 cup coffee or tea (without dairy products); 1 cup broth; 1 bowl gelatin

Snack: 1 glass fruit juice; 1 bowl gelatin

Lunch: 1 glass fruit juice; 1 glass water; 1 cup broth; 1 bowl gelatin

Dinner: 1 cup juice or water; 1 cup broth; 1 bowl gelatin; 1 cup of coffee or tea (without dairy products)